

Hector R Martinez Menchaca¹ and Gerardo Rivera Silva^{2*}

¹University of Louisville, Department of Orthodontics, Pediatric Dentistry and Special Care, USA.

²University of Monterrey, Health Science Division, Monterrey, NL, Mexico

Dates: Received: 18 June, 2016; Accepted: 08 July, 2016; Published: 09 July, 2016

***Corresponding author:** Dr. Gerardo Rivera Silva, PhD, Av. I. Morones Prieto # 4500 Pte, San Pedro Garza Garcia, NL, 66238, Mexico, Tel: +52 (81) 8215-1451; Fax: +52 (81) 8215-1000; E-mail: gerardo.rivera@udem.edu

www.peertechz.com

ISSN: 2394-8418

Clinical Image

Neonatal Teeth and Riga-Fede Disease




Figure 1:

Clinical Image

A 3 week-old male presented with ulceration on the ventral surface of the tongue. His parents informed that the patient had poor nutrients intake due to pain caused by tongue ulcer. The intraoral exploration showed ulceration of 10 mm diameter enclosed with a white fibrinous layer, situated on the ventral area of the tongue and two crows of neonatal teeth located in the mandibular anterior region (Figure 1). The incisal edges of teeth were smoothed about 2 mm. During a one-month follow-up his tongue ulcer disappeared.

Copyright: © 2016 Martinez Menchaca HR, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Martinez Menchaca HR, Silva GR (2016) Neonatal Teeth and Riga-Fede Disease. *J Dent Probl Solut* 3(1): 030-030. DOI: 10.17352/2394-8418.000030