

Xue Gao^{1,2}, Sha-Sha Huang¹, Yu Su¹,
Jin-Cao Xu and Pu Dai^{1,2*}

¹Department of Otorhinolaryngology, Head and Neck Surgery, PLA General Hospital, 28# Fuxing Road, Beijing, 100853, PR China

²Department of Otorhinolaryngology, General Hospital of the Rocket Force, 16# Xin Wai Da Jie, Beijing, 100088, PR China

Dates: Received: 28 June, 2016; Accepted: 07 July, 2016; Published: 08 July, 2016

*Corresponding author: Pu Dai, PhD, Department of Otorhinolaryngology, Head and Neck Surgery, PLA General Hospital, 28# Fuxing Road, Beijing, 100853, PR China, E-mail: daipu301@vip.sina.com

www.peertechz.com

Keywords: ARNSHL; Targeted deafness genes capture; Next-generation sequencing; TMC1

Research Article

Identification of Novel *TMC1* Compound Heterozygous Mutations Related to Autosomal Recessive Hearing Loss by Targeted Capture Sequencing

Abstract

Mutations in the transmembrane channel-like gene1 (*TMC1*) are known to cause autosomal dominant and recessive forms of nonsyndromic hearing loss DFNA36 and DFNB7/11, respectively. Here, we characterized a 5-year old girl with severe sensorineural hearing loss. By combining targeted capture of 140 known deafness genes, next-generation sequencing and bioinformatic analysis, we identified two compound heterozygous mutations *TMC1* c.1333C>T (p.R445C) and c.1765A>G (p.M589V) as the disease-causing mutations. Our results indicated that *TMC1* c.1333C>T (p.R445C) and c.1765A>G (p.M589V) lead to hearing loss in this family and highlight the power of combining deafness gene capture and next generation sequencing for identification of defective genes in sporadic cases with hearing loss. This approach will facilitate identification of the mutation spectrum of all known deaf genes.

Background

Hearing loss is a common sensory defect in humans. Nonsyndromic hereditary forms, in which hearing loss is the only clinical sign, are known to be genetically heterogeneous [1]. To date, more than 200 genetic loci and 140 responsible genes for hereditary hearing loss have been identified (Hereditary Hearing Loss Homepage, <http://hereditaryhearingloss.org/>). The extremely genetic heterogeneity of hearing impairment is a major challenge for traditional genetic testing and counseling. Targeted DNA capture and massively parallel sequencing are ideal tools to address this challenge.

Next generation sequencing, is a revolutionary technology that allows large amounts of genomic sequence information to be obtained rapidly and at a low cost [2]. Because of its large capacity to survey the whole exome and genome in an unbiased manner, NGS (next generation sequencing) is well suited to identifying the causative mutations of hereditary hearing loss. NGS has commonly been used to identify disease genes within even a limited number of patient samples [3-7]. Novel genes for non-syndromic [8,9] and syndromic [10] hearing loss have also been identified using the targeted NGS approach. Targeted deaf gene enrichment usually increases this proportion by at least 1000 fold [11,12]. In this study, we applied targeted capture of 140 known deafness genes, next-generation sequencing and bioinformatic analysis to analyze the genetic pathogenesis of one sporadic patient with severe sensorineural

TMC1 is predicted to encode membrane proteins with six transmembrane domains that may be involved in the functional maturation of cochlear hair cells [13]. Although the specific functions of the protein encoded by *TMC1* are unknown, bioinformatic analysis [14] and data obtained from *in vitro* heterologous systems

[13], suggest that *TMC1* functions as a membrane channel or transporter [15]. *TMC1* mutations cause nonsyndromic autosomal dominant and recessive hearing loss at the DFNA36 and DFNB7/11 loci, respectively. More than 30 recessive mutations in *TMC1* are associated with autosomal recessive nonsyndromic hearing loss (ARNSHL) at the DFNB7/11 locus in 39 families worldwide [14,16-24].

In this study, we identified one small ARNSHL pedigree with severe sensorineural hearing loss caused by the c.1333C>T (p.R445C) and c.1765A>G (p.M589V) mutation of *TMC1* through targeted deaf gene capture, next generation sequencing and bioinformatic analysis. This is the first study to identify *TMC1* c.1333C>T (p.R445C) and c.1765A>G (p.M589V) as the ARNSHL-associated mutation in this family.

Materials and Methods

Clinical data

Family C01701 is a two-generation Chinese family with autosomal recessive nonsyndromic hearing loss. To identify candidate mutations, DNA samples were obtained from 3 members of family C01701. Mutations of *GJB2* and *SLC26A4* had been excluded previously. Fully informed written consent was attained from two guardians. The study was approved by the Chinese PLA General Hospital ethics of research committees. Clinical information was gathered through multiple interviews with the family members. Medical history was obtained using a questionnaire regarding the following aspects of this condition: subjective degree of hearing loss, age at onset, evolution, symmetry of the hearing impairment, use of hearing aids, presence of tinnitus, medication, noise exposure, pathological changes in the

ear and other relevant clinical manifestations. Otoscopy, physical examination and pure tone audiometric examination (at frequencies from 250 to 8000 Hz) were performed to identify the phenotype. Immittance testing was applied to evaluate middle-ear pressure, ear canal volume and tympanic membrane mobility. Unaffected phenotype status was defined by a threshold lower than age- and gender-matched 50th percentile values for all frequencies measured. Physical examination of all members revealed no signs of systemic illness or dysmorphic features. CT scans of the temporal bone in the patient were performed. The diagnosis of profound sensorineural hearing impairment was made according to the WHO criteria based on audiometric examination. Tandem gait and Romberg tests were performed to evaluate balance.

Deafness gene capture and Illumina library preparation

Each DNA sample is quantified by agarose gel electrophoresis and Nanodrop (Thermo). Libraries were prepared using Illumina standard protocol. Briefly, 3 microgram of genomic DNA was fragmented by nebulization, the fragmented DNA is repaired, Illumina adapters are then ligated to the fragments, and the sample is size selected aiming for a 350–400 base pair product. The size selected product is PCR amplified (Each sample is tagged with a unique index during this procedure), and the final product is validated using the Agilent Bioanalyzer.

The amplified DNA was captured with a deafness related Gene Panel using biotinylated oligo-probes (MyGenostics GenCap Enrichment technologies). The probes were designed to tile along 140 deafness related genes. The capture experiment was conducted according to manufacturer's protocol. The enrichment libraries were sequenced on Illumina HiSeq 2000 sequencer for paired read 100bp.

Bioinformatics analysis

For sequencing analysis, high-quality reads were retrieved from raw reads by filtering out the low quality reads and adaptor sequences using the Solexa QA package and the cutadapt program (<http://code.google.com/p/cutadapt/>), respectively. SOAPaligner program was then used to align the clean read sequences to the human reference genome (hg19).

After the PCR duplicates were removed by the Picard software, the SNPs was firstly identified using the SOAPsnp program (<http://soap.genomics.org.cn/soapsnp.html>). Subsequently, we realigned the reads to the reference genome using BWA and identified the insertions or deletions (InDels) using the GATK program (http://www.broadinstitute.org/gsa/wiki/index.php/Home_Page). The identified SNPs and InDels were annotated using the Exome-assistant program (<http://122.228.158.106/exomeassistant>). Magic Viewer was used to view the short read alignment and validate the candidate SNPs and InDels. Nonsynonymous variants were evaluated by four algorithms, Ployphe, SIFT, PANTHER and Pmut, as described previously to determine pathogenicity.

Mutational analysis of *TMC1*

The segregation of the *TMC1* c.1333C>T (p.R445C) and c.1765A>G (p.M589V) mutations was tested in 3 family members

(I:1, I:2 and II:1), including the one whose gDNAs has been subjected to 140 deafness-associated gene NGS analysis, using PCR followed by bidirectional Sanger sequencing of the amplified fragments (ABI 3100, Applied Biosystems, USA). Nucleotide alteration(s) was identified by sequence alignment with the *TMC1* GenBank sequence using the Genetool software.

Multiple sequence alignment

Multiple sequence alignment was performed according to a Homologene program with default settings and the sequences NP_619636.2 (*H. sapiens*), XP_001093188.2 (*M. musculus*), XP_528322.2 (*P. troglodytes*), XP_002689695.2 (*B. taurus*), XP_005616877.1 (*C. lupus*), NP_083229.1 (*M. musculus*), NP_001101991.1 (*R. norvegicus*), NP_001006580.1 (*G. gallus*) and XP_695621.4 (*D. rerio*).

(http://www.ncbi.nlm.nih.gov/homologene?cmd=Retrieve&dopt=MultipleAlignment&list_uids=23670)

Results

Family and clinical evaluations

Family C01701 is a two-generation Chinese family with severe sensorineural hearing loss (Figure 1A) and includes 1 affected patients: II:1 (female, 5 years old). Hearing impairment was severe, prelingual and stable (Figure 1B).

Vestibular analysis was performed in the patient II:1 who did not complain about dizziness, vertigo, or imbalance. All position tests produced no nystagmus without vertigo sensation. II:1 did not have obvious delayed gross motor development. The physical examinations revealed no signs of systemic illness. High-resolution computed tomography of the temporal bone in II:1 was normal,




Figure 1: A. Pedigree of Family C01701 with ARNSHL Affected subject is denoted in black. Arrow indicates the proband; B. Audiogram of affected subject showed severe to profound sensorineural hearing loss. C. Electro-pherograms analysis of *TMC1* in family C01701 showing the compound heterozygous mutations (c.1333C>T and c.1765A>G) co-segregated with the phenotype.



References

- Friedman TB, Griffith AJ (2003) Human nonsyndromic sensorineural deafness. *Annu Rev Genomics Hum Genet* 4: 341-402.
- Metzker ML (2010) Sequencing technologies - the next generation. *Nat Rev Genet* 11: 31-46.
- Simpson DA, Clark GR, Alexander S, Silvestri G, Willoughby CE (2011) Molecular diagnosis for heterogeneous genetic diseases with targeted high-throughput DNA sequencing applied to retinitis pigmentosa. *J Med Genet* 48: 145-151.
- Kalay E, Yigit G, Aslan Y, Brown KE, Pohl E, et al. (2011) CEP152 is a genome maintenance protein disrupted in Seckel syndrome. *Nat Genet* 43: 23-26.
- Krawitz PM, Schweiger MR, Rodelsperger C, Marcelis C, Kolsch U, et al. (2010) Identity-by-descent filtering of exome sequence data identifies PIGV mutations in hyperphosphatasia mental retardation syndrome. *Nat Genet* 42: 827-829.
- Kuhlenbaumer G, Hullmann J, Appenzeller S (2011) Novel genomic techniques open new avenues in the analysis of monogenic disorders. *Hum Mutat* 32: 144-151.
- Musunuru K, Pirruccello JP, Do R, Peloso GM, Guiducci C, et al. (2010) Exome sequencing, ANGPL3 mutations, and familial combined hypolipidemia. *N Engl J Med* 363: 2220-2227.
- Rehman AU, Morell RJ, Belyantseva IA, Khan SY, Boger ET, et al. (2010) Targeted capture and next-generation sequencing identifies C9orf75, encoding taperin, as the mutated gene in nonsyndromic deafness DFNB79. *Am J Hum Genet* 86: 378-388.
- Walsh T, Shahin H, Elkan-Miller T, Lee MK, Thornton AM, et al. (2010) Whole exome sequencing and homozygosity mapping identify mutation in the cell polarity protein GPM2 as the cause of nonsyndromic hearing loss DFNB82. *Am J Hum Genet* 87: 90-94.
- Pierce SB, Walsh T, Chisholm KM, Lee MK, Thornton AM, et al. (2010) Mutations in the DBP-deficiency protein HSD17B4 cause ovarian dysgenesis, hearing loss, and ataxia of Perrault Syndrome. *Am J Hum Genet* 87: 282-288.
- Brownstein Z, Friedman LM, Shahin H, Oron-Karni V, Kol N, et al. (2011) Targeted genomic capture and massively parallel sequencing to identify genes for hereditary hearing loss in Middle Eastern families. *Genome Biol* 12: R89.
- Shearer AE, DeLuca AP, Hildebrand MS, Taylor KR, Gurrola J, et al. (2010) Comprehensive genetic testing for hereditary hearing loss using massively parallel sequencing. *Proc Natl Acad Sci U S A* 107: 21104-21109.
- Marcotti W, Erven A, Johnson SL, Steel KP, Kros CJ (2003) Tmc1 is necessary for normal functional maturation and survival of inner and outer hair cells in the mouse cochlea. *J Physiol* 574: 677-698.
- Kurima K, Peters LM, Yang Y, Riazuddin S, Ahmed ZM, et al. (2002) Dominant and recessive deafness caused by mutations of a novel gene, TMC1, required for cochlear hair-cell function. *Nat Genet* 30: 277-284.
- Vreugde S, Erven A, Kros CJ, Marcotti W, Fuchs H, et al. (2002) Beethoven, a mouse model for dominant, progressive hearing loss DFNA36. *Nat Genet* 30: 257-258.
- Hilgert N, Alasti F, Dieltjens N, Pawlik B, Wollnik B, et al. (2008) Mutation analysis of TMC1 identifies four new mutations and suggests an additional deafness gene at loci DFNA36 and DFNB7/11. *Clin Genet* 74: 223-232.
- Kalay E, Karaguzel A, Caylan R, Heister A, Cremers FP, et al. (2005) Four novel TMC1 (DFNB7/DFNB11) mutations in Turkish patients with congenital autosomal recessive nonsyndromic hearing loss. *Hum Mutat* 26: 591.
- Kitajiri SI, McNamara R, Makishima T, Husnain T, Zafar AU, et al. (2007) Identities, frequencies and origins of TMC1 mutations causing DFNB7/B11 deafness in Pakistan. *Clin Genet* 72: 546-550.
- Meyer CG, Gasmelseed NM, Mergani A, Magzoub MM, Muntau B, et al. (2005) Novel TMC1 structural and splice variants associated with congenital nonsyndromic deafness in a Sudanese pedigree. *Hum Mutat* 25: 100.
- Santos RL, Wajid M, Khan MN, McArthur N, Pham TL, et al. (2005) Novel sequence variants in the TMC1 gene in Pakistani families with autosomal recessive hearing impairment. *Hum Mutat* 26: 396.
- Tlili A, Rebeh IB, Aifa-Hmani M, Dhoubi H, Moalla J, et al. (2008) TMC1 but not TMC2 is responsible for autosomal recessive nonsyndromic hearing impairment in Tunisian families. *Audiol Neurootol* 13: 213-218.
- Gao X, Su Y, Guan LP, Yuan YY, Huang SS, et al. (2013) Novel compound heterozygous TMC1 mutations associated with autosomal recessive hearing loss in a Chinese family. *PLoS One* 8: e63026.
- Hassan MA, Shah AA, Szmida E, Smigiel R, Sasiadek MM, et al. (2015) A TMC1 (transmembrane channel-like 1) mutation (p.S320R) in a Polish family with hearing impairment. *J Appl Genet* 56: 311-316.
- Davoudi-Dehaghani E, Fallah MS, Tavakkoly-Bazzaz J, Bagherian H, Zeinali S (2015) Allelic heterogeneity among Iranian DFNB7/11 families: report of a new Iranian deaf family with TMC1 mutation identified by next-generation sequencing. *Acta Otolaryngol* 135: 125-129.
- Sirmaci A, Duman D, Ozturkmen-Akay H, Erbek S, Incesulu A, et al. (2009) Mutations in TMC1 contribute significantly to nonsyndromic autosomal recessive sensorineural hearing loss: a report of five novel mutations. *Int J Pediatr Otorhinolaryngol* 73: 699-705.
- Puente XS, Pinyol M, Quesada V, Conde L, Ordonez GR, et al. (2011) Whole-genome sequencing identifies recurrent mutations in chronic lymphocytic leukaemia. *Nature* 475: 101-105.
- Simpson CL, Justice CM, Krishnan M, Wojciechowski R, Sung H, et al. (2011) Old lessons learned anew: family-based methods for detecting genes responsible for quantitative and qualitative traits in the Genetic Analysis Workshop 17 mini-exome sequence data. *BMC Proc* 5: S83.
- Hanlon MR, Wallace BA (2002) Structure and function of voltage-dependent ion channel regulatory beta subunits. *Biochemistry* 41: 2886-2894.
- Keresztes G, Mutai H, Heller S (2003) TMC and EVER genes belong to a larger novel family, the TMC gene family encoding transmembrane proteins. *BMC Genomics* 4: 24.
- Li-Smerin Y, Hackos DH, Swartz KJ (2000) alpha-helical structural elements within the voltage-sensing domains of a K(+) channel. *J Gen Physiol* 115: 33-50.
- Labay V, Weichert RM, Makishima T, Griffith AJ (2010) Topology of transmembrane channel-like gene 1 protein. *Biochemistry* 49: 8592-8598.

Copyright: © 2016 Gao X, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Gao X, Huang SS, Su Y, Xu JC, Dai P (2016) Identification of Novel TMC1 Compound Heterozygous Mutations Related to Autosomal Recessive Hearing Loss by Targeted Capture Sequencing. *Scientific J Genetics Gen Ther* 2(1): 013-016.