

Luciano Silva*

University of Pernambuco, Brazilian Association of Odontology, Brazil

Dates: Received: 10 October, 2015; Accepted: 02 November, 2015; Published: 04 November, 2015

*Corresponding author: Luciano Silva, University of Pernambuco, Brazilian Association of Odontology - ABO - Recife, Tel: 55 81 30316824; E-mail: lucianobarreto2004@ig.com.br

www.peertechz.com

ISSN: 2455-5363

Keywords: Inflammation; Cytosine; Leukocytes

Review Article

A Literature Review of Inflammation and Its Relationship with the Oral Cavity

Abstract

Objective: This article presents a literature review which relates the processes, interactions, classification and the vascular and clinical characteristics of inflammation relating it with the oral cavity and describing the main cell types and chemical mediators used for its establishment.

Materials and Methods: it was used as source of research the following databases: MEDLINE, LILACS, BBO, Cochrane Library, Scielo, PubMed and the database of CAPES.

Results: much information was obtained concerning the chemical mediators as well as the main cell types responsible for the establishment of inflammation and its relationships with the oral cavity.

Introduction

The earliest known description of inflammation comes from the *Edwin Smith Papyrus*, one of the Egyptian papyri found in a tomb near Thebes. The *Papyrus* dates from about 1550 B.C., but it is undoubtedly a copy of ancient texts of the archaic period of Egyptian history (3200-2780 B.C.) Documents from Egyptian and other early civilizations leave no doubt that the features of inflammation were recognized from very early periods; although the understanding of the process took a long time to happen.

Inflammation can be considered as the first response of a living tissue to injury and may be referred to as the innate immunity, and may appear due to microbial infections, physical factors (trauma, radiation, temperature), chemical substances (irritant and corrosive chemicals), as well as tissue necrosis and hypersensitivity reactions. It is didactically characterized by the following quintet: redness (*rubor*), heat (*calor*), swelling (*tumor*), pain (*dolor*) and dysfunction of the organs involved (*functio laesa*). The first four characteristics were drawn up by Celsus nearly 2000 years ago; *functio laesa* was added to the definition of inflammation by Rudolf Virchow in 1858, considered the predecessor of modern pathology and social medicine. It was the latter who, in the field of inflammation, critically analyzed the meaning of the four key symptoms (redness, swelling, heat and pain) and postulated that inflammation could not be represented as a single process but rather constituted of various inflammatory processes.

Vascular and cellular considerations

A definition of inflammation is complicated, because the local vascular and tissue reactions often are accompanied by systemic effects that include malaise, fever, leukocytosis, metabolic disturbances and shock. The process of inflammation can be defined as the succession of changes which occurs in the living tissue when it is injured, provided that the injury is not of such a degree as at once to destroy its structure and vitality [1]. The inflammatory response consists of a vascular and

a cellular component. The vascular component represents changes in vessel diameter (vasodilation) and consequently increased blood flow (causing redness and heat), and increased vascular permeability resulting in loss of plasma into the tissue and formation of fluid exudate. The exudate contains many valuable proteins such as fibrin and immunoglobulin but it also gives rise to edema. The swelling distends the tissues; it may compress nerve endings and thus cause pain.

According to its duration, inflammation is described as acute or chronic. Sometimes the acute process subsides but the stimulus persists sufficiently to evoke a subsequent chronic inflammation. In other cases, with a stimulus that typically induces chronic inflammation, the tissue response may be acute in type for the first day or so. Suffice it to say at this stage that the tissue response differs considerably in acute and chronic. The cellular component involves the movement of white blood cells (leukocytes) from the blood vessels into the inflamed tissue. They extravasate from the capillaries into tissue, and act as phagocytes, picking up bacteria and cellular debris. They may also aid by walling off an infection and preventing its spread. Influx of neutrophils is one of the earliest stages of the inflammatory response. These cells mount a rapid, non-specific phagocytic response. Later, monocytes/macrophages and cells of other lymphocyte lineages (specific subsets of T cells and B cells) appear at the site of injury; cell types which are associated with antigen-specific and more tightly regulate immune responses and once activated also produce protective and inflammatory molecules.

The lymphocytes are the primary cells of the immunologic system, and have developed one of the more sophisticated intelligent defense mechanisms in the biological system. The review of ALAM and GORSKA [2], enlightens that the T lymphocytes play a major role in the organization of the immune response, eliminating intracellular pathogens (virus, and bacteria) through the generation of cytotoxic T lymphocytes. The B lymphocytes protect the organism against extracellular pathogens by the production of antibodies. The

natural killer cells (NK) are an important component of the innate immunity. The dendritic B cells begin the immunologic response by presenting antigens to the T lymphocytes. The authors resume that the interaction among the T lymphocytes, B lymphocytes, dendritic cells and the natural killer cells (NK) constitutes the fundamental defensive mechanism of the host.

The mechanism against pathogenic microbes demands different responses depending on the characteristic of the pathogen and on the attacked tissue. CHAPLIN [3], claimed that the organism of the host has developed innate and adaptive immune mechanisms of defense, being the former unspecific, attacking any nonself structure or antigen, and the latter, on the other hand, extremely specific. Both types of immune response act together in order to eliminate pathogenic antigens through the discrimination of what is self and what is not.

The changes described in blood flow, increased permeability of blood vessels and escape of cells from the blood into the tissues (diapedesis) are basically the same whatever the cause and wherever the site of inflammation is. Thermal injuries cause permanent occlusion of the blood vessels in a narrow zone of tissue immediately surrounding the burned site. Beyond the zone where the injury is less intense, blood flow is reestablished after a short initial period of vasoconstriction. Immediately after injury, there is transient ischemia due to vasoconstriction, particularly of the arterioles. The sequelae of acute inflammation depend upon the type of tissue involved and the amount of tissue destruction, which depend, in turn, upon the nature of the injurious agent. The possible outcomes of acute inflammation can be either healing or chronic inflammation. Chronic inflammation is characterized by a dominating presence of macrophages in the injured tissue. These cells provide a powerful defensive mechanism in the body, but the mediators they release are injurious to the organism's own tissues, as well as invading agents. This is why chronic inflammation is almost always accompanied by tissue destruction [4].

Inflammation is referred to, didactically, as *acute* when it lasts two days or three weeks and as *chronic* when more prolonged. The typical case of acute inflammation is characterized by vasodilatation, by exudation of plasma and the emigration of neutrophilic leucocytes into the injured tissue. But not all the examples of acute inflammation exhibit neutrophilic infiltration and, conversely, neutrophils may be associated with prolonged and therefore *chronic* inflammation. Between acute and chronic there occurs a wide range of overlapping processes and those at halfway point are sometimes referred to as *subacute* inflammation. The main events in acute inflammation are vascular in origin and remarkably consistent for a wide range of stimuli. Nevertheless, the response exhibits considerable differences that depend, first, on factors related to the injury or infection and, second, on the condition of the host and nature of the tissue involved [5]. On the other hand, inflammation is said to be chronic when its duration is prolonged for months or years, or even indefinitely. Its prolonged course is due to persistence of the causative factor in the tissues, whether the factor be infection or inanimate foreign body. The tissues committed are infiltrated by macrophages, lymphocytes as well as plasma cells. In addition, there is a varying but often form substantial proliferation of fibroblasts forming collagen, blood vessels and of tissue histiocytes.

The role of cytokines

The immune system is a complex organ with both professional and non-professional cells involved, whose main objective is to allow the most developed organisms to endure in a world where microorganisms try to make use of them to survive. In short; it is a sophisticated and interconnected system that makes human existence possible; and so communication between its main components is a major issue. Therefore, the competent cells of the immune system make use of substances that may be unleashed in the blood stream to signalize the need of the cell recruitment or the production of specific substances. It is accomplished by direct cell-cell contacts, by contacts with non-self-factors; like pathogen compounds, and by soluble proteins. This immune response is obtained partially by the synchronized synthesis and function of the small cytokines called chemokines (CKs) and their action on their counterparts, the chemokine receptors (CKRs); without which the cells would never respond. In fact, CKRs define the subpopulation that will be recruited in each step of the immune responses. The name chemokines is derived from chemoattractant cytokines, and is related to their capacity of inducing directed chemo taxis in nearby responsive cells.

The cytokine family is large with still unanswered questions. It includes the Interleukins (IL), Interferon (IFN), Tumor Necrosis Factor (TNF), Colony-stimulating Factor (CSF), Chemokines (CKs), and Growth Factor (GF) [3].

Interleukins – ranging from IL1 to IL32 - are different among each other, have different functions and are secreted by different cells. Interferon is subdivided in IFN α (leukocyte IFN - inhibitor of viral replication) IFN β (fibroblast IFN – inhibitor of viral replication) and IFN γ (Secreted by lymphocytes – immune regulatory functions). Tumor Necrosis Factor is represented by TNF α (secreted frequently by monocytes) and TNF β (secreted by T cells). Colony-stimulating Factor (G-CSF, M-CSF, GM-CSF and others) – able to make bone-marrow cells differentiation into specific cell types, like monocytes, macrophages and neutrophils. Chemokines (Eotaxin and many others) are very important in controlling the migration of cells between and within tissues. Growth Factor (IGF, TGF and many others) was originally identified because of non-immune-related functions but may have effects on immune cells. Some of them are considered pro-inflammatory and can be induced during an immune response to promote cells of the immune system to a site of infection, whereas others are considered homeostatic and are involved in controlling the migration of cells during normal processes of tissue maintenance or development. The first cytokine identified to have chemotactic activity was interleukin-8 (IL-8), which proved to be a selective neutrophil chemoattractant [6].

Not only is the inflammatory bone destruction governed by local cytokine production but also the normal bone remodeling throughout the lifetime of an individual. Some of the factors which regulate physiologic bone remodeling are prostaglandins (PGs), IL-1, TNF- α and possibly IL-6 [7].

The production of cytokines at the sites of inflammation in the oral tissues is part of the host response which is essentially protective in nature. Both under-activity (hypo-responsiveness)

and over-activity (hyper-responsiveness) of the host response, such as unrestricted production of cytokine(s), may result in tissue destruction. Traditionally, the immune functions have been thought to be regulated by signals originating within the immune system. It is now evident that the immune system is regulated in part by the central nervous system (CNS), acting principally via the hypothalamic-pituitary adrenal (HPA) axis and the sympathetic nervous system (SNS) [8,9]. The pathways between immune system and brain seem to be bidirectional and the aim is maintenance of homeostasis. The SNS provides a major integrative and regulatory pathway for this communication. The sympathetic innervation of lymphoid tissue, the presence of adrenergic receptors on immune cells (B and T lymphocytes, macrophages) and studies of catecholamine interactions with the immune system [10], provide substantial evidence for the role of SNS in immune regulation. Furthermore, the cellular products of an activated immune system, namely cytokines, can signal to the brain. The cytokines TNF- α , IL-6 and IL-1 seem to be involved in the cross-talk between brain and immune system via corticotropin-releasing hormone (CRH) secretion and hence, they activate both the HPA and the SNS [11,12].

The development of methods for the study of leukocyte migration *in vitro* [13], facilitated the discovery of several chemoattractants, such as complement fragments C3a and C5a, arachidonic acid derivatives such as leukotriene B₄ (LTB₄) and 12-hydroxy-eicosanotetraenoic acid (12-HETE), and platelet activating factor (PAF) [14].

Inflammation in the oral cavity

The oral cavity is exposed in many ways to a number of potential microbial agents firstly because of its main functions: nourishment and communication. Other than these factors, some treatments applied on the teeth may favor the deposition of dental plaque associated with extra strength applied on the teeth, like orthodontics. Even more, inflammation in orthodontics also comes from the forces applied on the teeth, and the presence of osteoclasts at location of alveolar bone remodeling is antecedent to orthodontic tooth movement. Cell recruitment and clearance are the mechanisms by which osteoclast population are regulated [15]. During orthodontic force application, the periodontal ligament (PDL) undergoes hyalinization, a light microscope term describing the loss of cells from an area of the ligament because of necrosis. This tissue damage prevents the tooth from moving until the adjacent bone and the necrotic tissue are removed by osteoclasts. As a result, the tooth movement reaches a plateau which lasts about 10 to 20 days in humans [16,17].

Periodontitis and pulpitis are the most common infections in odontology and are indeed commonly found in the daily practice in surgery offices. Periodontitis shares many pathologic features with other inflammatory diseases with concomitant bone resorption such as rheumatoid arthritis (RA), with accumulative evidence that both conditions manifest as a result of an imbalance between pro-inflammatory and anti-inflammatory cytokines [18]. The increased local production of pro-inflammatory cytokines either directly enhances the proliferation and/or activity of cells in the osteoclast lineage or indirectly affects the production of the essential osteoclast differentiation factor, receptor activator of NF- κ B ligand, and/or its

soluble decoy receptor, osteoprotegerin, by osteoblast/stromal cells [19].

As for what concerns root resorptions; one study showed that only interleukin-1 α has a potent capacity, to increase it [20].

ESSAYAN, 1998 studied the role of the IL-1 cytokine family (interleukin-1), which represents a group of proteins that possesses synergic and contrasting biological responses. In agreement with their study, IL-1 and its precursory forms are strongly involved in the establishment of inflammation and host defense [21]. The continuous exposure to many sorts of biological, physical and chemical agents makes the mouth more prone to inflammation and infections, mainly caused by bacterial influence. Its main tissues are, in normal conditions, well adapted to endure; supporting even abrupt thermal changes, but sometimes, as a result of low immunity or high virulence of a determined agent (or even a combination of both), it may develop an inflamed site elsewhere in the mouth, where varieties of anaerobic gram negative bacteria exist and the LPS (lipopolysaccharide – a bacterial pathogenic element) of such bacteria is thought to be a virulence factor which plays a role in the development of dental diseases [22-24]. Once inflammation such as pulpitis, gingivitis, or periodontitis has been established, an inflammatory infiltrate is formed consisting of different cell types such as neutrophils, macrophages and lymphocytes which will produce different cytokine subtypes responsible for the immunopathology of the diseases.

Chronic marginal gingivitis is characterized clinically by gingival redness, edema, bleeding, changes in contour, loss of tissue adaptation to the teeth, and increased flow of gingival crevicular fluid (GCF) [25,26]. Its origins is usually associated with dental plaque which irritates the surrounding tissues because of the release of toxins produced by microorganisms and the natural course of the disease may many times be modified by the action of hormones. Pregnancy is accompanied by an increase in the levels of both progesterone and estrogen which, by the third trimester, reaches levels 10-30 times than seen during the typical menstrual cycle. The changes in the levels of hormone lead to the conclusion that female steroid hormones may have dual effects on the pathogenesis of pyogenic granuloma in pregnancy.

Development of gingivitis necessarily requires the presence of plaque bacteria [27,28] which is believed to cause pathological changes in the tissues by both direct and indirect means [29]. Histopathologic observations have led to the subdivision of gingivitis into 3 stages, which are divided didactically for the understanding of the process. The first stage usually begins with an initial lesion which appears 2 or 4 days after initial injury, where there is a transitory period of vasoconstriction followed by vasodilatation and consequent increase of blood flow that brings along neutrophils (PMNs), progressively adhering to the endothelial walls getting ready for diapedesis. Vascular changes and collagen degradation represent the initial changes probably due to chemotactic attraction of neutrophils to the contaminated area by direct vasodilatory effects of bacterial products, as well as activation of host systems such as the complement and kinin systems and arachidonic acid pathways [30]. The second stage is a continuation of the process represented by the persistence of the

acute inflammatory response. Capillaries engorge and proliferate as a result of blood congestion, which increases the permeability and the inflow of plasma and the diapedesis, with a mixture of the first lineage of immune cells, which is unspecific, and the adaptive immunity, with the lymphocytes as their main representatives, preparing the area for a transition from acute to chronic. And finally the third stage would be the establishment of chronic gingivitis characterized by slow venous return and color change from red to blue hue and the intense population of lymphocytes B and the formation of granulation tissue rather than collagen bundles [31].

In situ production of pro-inflammatory cytokines is generally seen in inflamed pulp and/or gingiva as part of the inflammatory response [32,33]. The clinical signs and the discomfort of gingivitis have been attributed to pathological tissue changes which commit gingival microcirculation [34], a mechanism which involves capillaries and small venules right at the site of injury and the adjacent arterioles. Bacterial LPS and cytokines have only been implicated in the vascular component of edema formation [35], however their possible effect on the interstitium has not yet been explored. Following the expectable path of untreated gingivitis, it can progress to a more severe stage when the inflammatory process extends to the periodontal ligament and alveolar bone, compromising directly the anchoring apparatus of the teeth [36].

On the other hand, in periodontitis, bone resorption is mediated largely by the increased local production of pro-inflammatory cytokines, such as IL-1 α , IL-1 β , and TNF- α . The LPS released from the periodontopathic bacteria triggers the synthesis of IL-1 and TNF- α from macrophages. IL-1 and TNF- α stimulate the production of a variety of other inflammatory mediators (cytokines, nitric oxide and PGE2), matrix-degrading enzymes (metalloproteinases), and decrease the synthesis of glycosaminoglycans, enhancing bone destruction [37,38].

The degradation and removal of hard tissue are a cellular event brought about by giant multinucleated cells formed through asynchronous fusion of mononuclear cells belonging to the macrophage lineage and originating from the hematopoietic system [39]. They are called clasts and are easy to identify under the light microscope because of their size (50 to 100 μ m), their multinucleation (2 to 10 nuclei per cell), and their association with the surface of bone (occupying shallow depressions known as Howship's lacunae). Such cells are also intermediated by cytokines who induce and stimulate their recruitment.

The field of odontology which deals with bone resorption and neoformation is orthodontics, for the fact that its basis is the application of mechanical forces which produce organized periodontal tissue remodeling generating tooth movement, with forces transmitted from the dental roots to the periodontium where cells are stimulated to remodel the surrounding matrices. The forces applied will necessarily cause an inflamed area in the surrounding tissues of a tooth, causing bone resorption in the pressure sites and bone deposition in the tension sites [40-43]. There seems to be a role of cytokines such as IL-1 α , IL-1 β , and TNF- α in the process [44,45].

As previously mentioned, several infectious, autoimmune,

but also allergic and neoplastic diseases have been attributed to the imbalance between pro-inflammatory and anti-inflammatory cytokines and therefore, the regulation of pro/anti-inflammatory cytokines has attracted considerable interest [46]. Components of the innate immunity such as antigen presenting cells (APCs), dendritic cells, and monocytes/macrophages, as well as components of the acquired (adaptive) immunity such as T helper (Th) lymphocytes, are all involved in immunoregulation.

To mediate an effective response, leukocytes must find their way to sites of infection or inflammation. Leukocyte invasion of tissues can be induced by chemotactic activity of several substances—including interleukin-1 (IL-1), tumor necrosis factor- (TNF- α), and bacterial lipopolysaccharide (LPS)—that cause leukocyte emigration when injected *in vivo* [47]. All such compounds induce the production of chemoattractants, which in turn cause leukocyte migration. Therefore, chemotactic activity includes the receptor-mediated gradient perception and must be measured by the ability of a chemoattractant to induce directed leukocyte migration *in vitro* [47].

The antigen presenting cells (APC) are those that express high levels of class II MHC molecules; with internalizing ability to process and expose foreign antigens in the fitting MHC. Therefore, the immune response comes mainly from the action of four cellular types: T cells (fundamental in the elimination of intracellular pathogens such as virus and bacteria) through the generation of cytotoxic T cells. B cells, which defend the organism against extracellular pathogens by the production of antibodies. Natural killer cells (NK), which constitute a subseries of positive cytotoxic lymphocytes CD56, which, in spite of belonging to the innate immunologic system for not presenting surface receptors, are fundamental for the recognition of pathogens, and Dendritic cells (DC), cells, which activate the immunologic response by the exhibition and consequent presentation of foreign antigens to the T cells [48,49].

Th lymphocytes are divided into two subclasses; Th1 and Th2. Th1 cells primarily secrete IFN- γ , and IL-2 which enhance cellular immunity, whereas Th2 cells secrete a different set of cytokines, primarily IL-4, IL-10, IL-13 and IL-9, which enhance humoral immunity [50,51]. The antigen -inexperienced (naïve) CD4+ Th0 cells can differentiate to either Th1 or Th2 and the differentiation is strongly dependent on the cytokines produced by cells of the innate immune system [51]. IL-12 produced by activated monocytes/macrophages, represents a major inducer of Th1 differentiation and hence cellular immunity. IL-12, together with TNF- α and IFN- γ act synergistically in inflammation and further promote Th1 responses and hence considered major pro-inflammatory cytokines [51]. Th1 and Th2 responses are mutually inhibitory. Thus IL-12 and IFN- γ inhibit Th2 and vice versa, IL-4 and IL-10 inhibit Th1 responses and the production of pro-inflammatory cytokines. IL-4 and IL-10 are the major anti-inflammatory cytokines and an increasing body of evidence suggests that catecholamines inhibit selectively the Th1 functions and pro-inflammatory cytokines and favor the Th2 responses and anti-inflammatory cytokines [8].

Cytokines and chemokines are redundant secreted proteins with growth, differentiation, and activation functions that regulate and

determine the nature of immune responses and control immune cell trafficking and the cellular arrangement of immune organs. Which cytokines are produced in response to an immune insult determines initially whether an immune response develops and subsequently whether such response is cytotoxic, humoral, cell-mediated, or allergic [52]. Cytokines act on cells through transmembrane cell surface receptors. Binding of the cytokine to the receptor elicits its cellular response by activating an intracellular signal transduction pathway that ultimately leads to induction of new gene transcription and synthesis of new cellular proteins. Most cytokine receptors signal using one of the Janus Kinase (Jak) family of proteins. Specific Jak proteins associate with the cytoplasmic domains of cytokine receptors. When binding of the cytokine activates the receptor, the Jak phosphorylates its respective STAT protein, causing the STAT to dimerize and translocate into the nucleus, where it initiates new gene transcription [3].

Increased vascular permeability can be induced by cytokines such as IL-1 and TNF- α acting either directly at the endothelial cells or indirectly by inducing other inflammatory mediators that affect the vascular endothelium [53,54]. In addition, LPS can increase vascular permeability not only through induction of pro-inflammatory cytokines, but also affecting directly the endothelial barrier [55,56].

The *periapical lesion* represents a product of the body's immune response to bacterial infection of the pulp, with the purpose to wall-off the infection within the confines of the root canal system [57]. In many respects this periapical inflammatory response recapitulates the pulpal response to infection, with the additional feature of periapical bone destruction. A mixed infiltrate of T and B lymphocytes, polymorphonuclear leukocytes (PMNs), macrophages, plasma cells, natural killer (NK) cells, eosinophils and plasma cells has been characterized [58-63]. NK cells are thought to represent a third lineage of lymphoid cells which, when activated, have the morphology of a large granular lymphocyte [3].

There are controversial results regarding the predominant infiltrating cell type in periapical lesions. Such periapical alterations are didactically postulated as chronic lesions with either lymphocytes [64,65], or macrophages [61,63] generally reported to be most numerous. T cells consistently outnumber B cells [65,66]. Of the T lymphocytes, both Th and T suppressor cells have been identified with temporal differences regarding their relative concentration [63,66]. The variety of inflammatory cells reveals that both non-specific and specific immune responses are mediated in the pulp and periapical tissues in response to infection. Therefore, the periapical lesion provides a suitable model for studying local immune responses. The superior cervical ganglion (SCG) supplies with sympathetic innervation the oral tissues [67,68]. By removal of the SCG, the local control of immune responses by sympathetic nerves can be studied in periapical lesions.

The interstitial fluid (IF) represents extracellular fluid located outside the blood vessels. Interstitial fluid bathes the cells of the tissues and it provides a mean of Delivering nutrients to the cells, intercellular communication and removal of metabolic waste. Plasma, the major component of blood, communicates with the

interstitial fluid through pores and intercellular clefts in the capillary endothelium. Under physiologic conditions, the endothelium of the capillaries and proximal portion of the venules forms a semipermeable membrane that permits free movement of water and small molecules to and fro across the endothelium but normally restricts the passages of plasma protein. The capillary wall acts as a filtration "barrier". Fluid equilibrium across this endothelial barrier is maintained by the hydrostatic pressure of the capillary blood being balanced by the equal and opposite restraint of the osmotic pressure of the plasma proteins. Most of the fluid within the capillaries is retained, but some fluid filters normally through pores between the cells, pushed by the pressure difference between the capillary blood and the IF. Water and small solutes can pass freely through these pores. The net effect of the hydrostatic pressure alone is a net loss of water and solutes from plasma to the IF. The capillary wall is, however, nearly impermeable to the plasma proteins and lipids but following inflammation, the capillary permeability for proteins increases. Because the capillary wall is permeable to water, but essentially impermeable to the plasma proteins, these molecules generate a colloid osmotic pressure gradient that draws water out of the interstitium and into the plasma. This pressure is proportional to the difference in protein concentration between the plasma and the IF.

Nevertheless, there is little information available regarding intra- and extra-vascular fluid volumes in dental tissues (oral mucosa, pulp) during health and disease. The absence of such information from the dental pulp is mainly due to the nature of the tissue that does not allow easy access to the interstitium. Its inextensible walls creates a hermetic cavity inside which blood runs bringing in nutrients and defense cells, while removes metabolic wastes from inside the pulp. However, attempts have been made to isolate dentinal fluid, which is a pulpal IF that moves across the dentinal tubules, in order to measure COP and protein concentration [69-71]. In order to collect dentinal fluid, it is mandatory to drill a deep cavity in the dentin and the consequent removal of the smear layer. These are rather invasive procedures that may affect the pulpal blood flow and subsequently the rate and direction of dentinal fluid flow [72,73]. The radius of the exposed tubules [74], and the disruption of the odontoblast layer as a result of the preparation trauma [75], may yet influence the dentinal fluid composition regarding plasma proteins. It is therefore evident that there is a need for a less invasive and traumatic method for the due isolation of pulp IF in order to explore its microenvironment.

Conclusions

Inflammation is a complex process and a necessary defense strategy in search of the reestablishment of health. Its classification, cell types involved, cytokines and chemokines and other substances give us a hint of its nature and importance, although much more is to be discovered, mostly as for what concerns the oral cavity.

References

1. Adami JG (1909) Inflammation: an introduction to the study of pathology. London.
2. Alam R, Gorska M (2003) Lymphocytes. Allergy and Clinic. Immunol 111:
3. Chaplin DD (2010) Overview of the immunologic response. J Allergy Clin Immunol 125: S3-23.

4. Trowbridge HO, Emling RC (1997) *Inflammation. A Review of the Process* (5th ed). Quintessence Publishing 200.
5. Robbins SL (1968) *Textbook of pathology*, ed. 3, Philadelphia, W.B. Saunders Co.
6. Yoshimura T, Matsushima K, Tanaka S, Robinson EA, Appella E, et al. (1987) Purification of a human monocyte-derived neutrophil chemotactic factor that has peptide sequence similarity to other host defense cytokines. *Proc Natl Acad Sci USA* 84: 9233-9237.
7. Rodan GA (1992) Introduction to bone biology. *Bone* 13: S3-6.
8. Elenkov IJ, Wilder RL, Chrousos GP, Vizi ES (2000) The sympathetic nerve--an integrative interface between two supersystems: the brain and the immune system. *Pharmacol Rev* 52: 595-638.
9. Haddad JJ, Saadé NE, Safieh-Garabedian B (2002) Cytokines and neuro-immune-endocrine interactions: a role for the hypothalamic-pituitary-adrenal revolving axis. *J Neuroimmunol* 133: 1-19.
10. Madden KS, Sanders VM, Felten DL (1995) Catecholamine influences and sympathetic neural modulation of immune responsiveness. *Annu Rev Pharmacol Toxicol* 35: 417-448.
11. Del Rey A, Besedovsky H, Sorkin E, Dinarello CA (1987) Interleukin-1 and glucocorticoid hormones integrate an immunoregulatory feedback circuit. *Ann N Y Acad Sci* 496: 85-90.
12. Chrousos GP (1995) The hypothalamic-pituitary-adrenal axis and immune-mediated inflammation. *N Engl J Med* 332: 1351-1362.
13. Boyden S (1962) The chemotactic effect of mixtures of antibody and antigen on polymorphonuclear leucocytes. *J Exp Med* 115:453-466.
14. Schroder JM (2000) Chemoattractants as mediators of neutrophilic tissue recruitment. *Clin Dermatol* 18: 245-263.
15. Noxon SJ, King GJ, Gu G, Huang G (2001) Osteoclast clearance from periodontal tissues during orthodontic tooth movement. *J. Am. Orthod. Dentofacial Orthop* 120: 466-476.
16. Rygh P (1974) Elimination of hyalinized periodontal tissues associated with orthodontic tooth movement. *Scand J Dent Res* 82: 57-73.
17. Hixon EH, Aasen TO, Clark RA, Arango J, Klosterman R, et al. (1970) On force and tooth movement. *Am J Orthod* 57: 476-489.
18. Bartold PM, Marshall RI, Haynes DR (2005) Periodontitis and rheumatoid arthritis: a review. *J Periodontol* 76: 2066-2074.
19. Boyce BF, Li P, Yao Z (2005) TNF-alpha and pathologic bone resorption. *Keio J Med* 54: 127-131.
20. Rivollier A, Mazzorana M, Tebib J, Piperno M, Aitsiselmi T, et al. (2004) Immature dendritic cell transdifferentiation into osteoclasts: a novel pathway sustained by the rheumatoid arthritis microenvironment. *Blood* 104: 4029-4037.
21. Silva LB, Guimaraes CS, Santos RA (2008) Immunology of root resorption: a literature review. *Indian J Dent Res* 19: 340-343.
22. Slots J, Genko RJ (1984) Black-pigmented *Bacteroides* species, *Capnocytophaga* species, and *Actinobacillus actinomycetemcomitans* in human periodontal disease: virulence factors in colonization, survival, and tissue destruction. *J Dent Res* 63: 412-421.
23. Warfvinge J, Dahlen G, Bergenholtz G (1985) Dental pulp response to bacterial cell wall material. *J Dent Res* 64: 1046-1050.
24. Larjava H, Uitto VJ, Eerola E, Haapasalo M (1987) Inhibition of gingival fibroblast growth by *Bacteroides gingivalis*. *Infect Immun* 55: 201-205.
25. Cimasoni G (1983) Crevicular fluid updated. *Monogr Oral Sci* 12: III-VII 1-152.
26. Greenstein G (1984) The role of bleeding upon probing in the diagnosis of periodontal disease. A literature review. *J Periodontol* 55: 684-688.
27. Løe H, Theilade E, Jansen SB (1965) Experimental gingivitis in man. *J Periodontol* 36: 177-187.
28. Theilade E, Wright WH, Jensen SB, Løe H (1966) Experimental gingivitis in man. II. A longitudinal clinical and bacteriological investigation. *J Periodont Res* 1: 1-13.
29. Page R (1986) Gingivitis. *J Clin Periodontol* 13: 345-359.
30. Attström R, Egelberg J (1970) Emigration of blood neutrophils and monocytes into the gingival crevices. *J Periodontol Res* 5: 48-55.
31. Payne WA, Page RC, Ogilvie AL, Hall WB (1975) Histopathologic features of the initial and early stages of experimental gingivitis in man. *J Periodont Res* 10: 51-64.
32. Van Dyke TE, Lester MA, Shapira L (1993) The role of the host response in periodontal disease progression: implications for future treatment strategies. *J Periodontol* 64: 792-806.
33. Zehnder M, Delaleu N, Du Y, Bickel M (2003) Cytokine gene expression--part of host defence in pulpitis. *Cytokine* 22: 84-88.
34. Del Fabbro M, Francetti L, Bulfamante G, Cribiù M, Miserocchi G, et al. (2001) Fluid dynamics of gingival tissues in transition from physiological condition to inflammation. *J Periodontol* 72: 65-73.
35. Daffonchio L, Novellini R, Bertuglia S (2002) Protective effect of ketoprofen lysine salt on interleukin-1beta and bradykinin induced inflammatory changes in hamster cheek pouch microcirculation. *Inflamm Res* 51: 223-228.
36. Kornman KS (1986) The role of supragingival plaque in the prevention and treatment of periodontal diseases. A review of current concepts. *J Periodontol Res* 21: 5-22.
37. Meyer O (2003) Role of TNF-alpha and cytokines in the pathophysiology of rheumatoid arthritis. Therapeutic perspectives. *Bull Acad Natl Med* 187: 935-954.
38. Hong CY, Lin SK, Kok SH, Cheng SJ, Lee MS, et al. (2004) The role of lipopolysaccharide in infectious bone resorption of periapical lesion. *J Oral Pathol Med* 33: 162-169.
39. Silva LB, Guimaraes CS, Santos RA (2008) Immunology of root resorption: a literature review. *Indian J Dent Res* 19: 340-343.
40. Reitan K (1954) Parodontal reconstruction in the framework of orthodontics. *Zahnartzl Welt* 9: 2/35.
41. Rygh P (1973) Ultrastructural changes in pressure zones of human periodontium incident to orthodontic tooth movement. *Acta Odontol Scand* 31: 109-122.
42. Rygh P (1976) Ultrastructural changes in tension zones of rat molar periodontium incident to orthodontic tooth movement. *Am J Orthod* 70: 269-281.
43. Brudvik P, Rygh P (1993) The initial phase of orthodontic root resorption incident to local compression of the periodontal ligament. *Eur. J. Orthod* 15: 249-263.
44. Davidovitch Z, Nicolay OF, Ngan PW, Shanfeld JL (1998) Neurotransmitters, cytokines, and the control of alveolar bone remodeling in orthodontics. *Dent Clin North Am* 32: 411-435.
45. Saito M, Saito S, Ngan PW, Shanfeld J, Davidovitch Z (1991) Interleukin 1 beta and prostaglandin E are involved in the response of periodontal cells to mechanical stress in vivo and in vitro. *Am J Orthod Dentofacial Orthop* 99: 226-240.
46. Elenkov IJ, Chrousos GP (1999) Stress Hormones, Th1/Th2 patterns, Pro/ Anti-inflammatory Cytokines and Susceptibility to Disease. *Trends Endocrinol Metab* 10: 359-368.
47. Silva TA, Garlet GP, Fukada SY, Silva JS, Cunha FQ (2007) Chemokines in Oral Inflammatory Diseases: Apical Periodontitis and Periodontal Disease. *JDR* 86: 306-319.
48. Münz C, Dao T, Ferlazzo G, de Cos MA, Goodman K, et al. (2005) Mature myeloid dendritic cell subsets have distinct roles in activation and viability of circulating human natural killer cells. *Blood* 105: 266-273.

49. Osugi Y, Vuckovic S, Hart DN (2002) Myeloid blood cd11c+ dendritic cells and monocyte-derived dendritic cells differ in their ability to stimulate t lymphocytes. *Blood* 100: 2858-2866.
50. Abbas AK, Murphy KM, Sher A (1996) Functional diversity of helper T lymphocytes. *Nature* 383: 787-793.
51. Fearon DT, Locksley RM (1996) The instructive role of innate immunity in the acquired immune response. *Science* 272: 50-53.
52. Borish LC, Streinke JW (2003) 2 Cytokines and chemokines. *J Allergy Clin Immunol* 111:460-475.
53. Rossi V, Breviario F, Ghezzi P, Dejana E, Mantovani A (1985) Prostacyclin synthesis induced in vascular cells by interleukin-1. *Science* 229: 174-176.
54. Bussolino F, Breviario F, Tetta C, Aglietta M, Mantovani A, et al. (1986) Interleukin 1 stimulates platelet-activating factor production in cultured human endothelial cells. *J clin Invest* 77: 2027-2033.
55. Bannerman DD, Goldblum SE (1999) Direct effects of endotoxin on the endothelium: barrier function and injury. *Lab Invest* 79: 1181-1199.
56. Bannerman DD, Goldblum SE (2003) Mechanisms of bacterial lipopolysaccharide-induced endothelial apoptosis. *Am J Physiol Lung Cell Mol Physiol* 284: L899-914.
57. Stashenko P (1990) Role of immune cytokines in the pathogenesis of periapical lesions. *Endod Dent Traumatol* 6: 89-96.
58. Cymerman JJ, Cymerman DH, Walters J, Nevins AJ (1984) Human T lymphocyte subpopulations in chronic periapical lesions. *J Endod* 10: 9-11.
59. Torabinejad M, Kettering JD (1985) Identification and relative concentration of B and T lymphocytes in human chronic periapical lesions. *J Endod* 11: 122-125.
60. Gao Z, Mackenzie IC, Rittman BR, Korszun AK, Williams DM, et al. (1998) Immunocytochemical examination of immune cells in periapical granulomata and odontogenic cysts. *J Oral Pathol* 17: 84-90.
61. Kopp W, Schwarting R (1989) Differentiation of T lymphocyte subpopulations, macrophages, and HLA-DR-restricted cells of apical granulation tissue. *J Endod* 15: 72-75.
62. Stashenko P, Yu SM (1989) T helper and T suppressor cell reversal during the development of induced rat periapical lesions. *J Dent Res* 68: 830-834.
63. Kawashima N1, Okiji T, Kosaka T, Suda H (1996) Kinetics of macrophages and lymphoid cells during the development of experimentally induced periapical lesions in rat molars: a quantitative immunohistochemical study. *J Endod* 22: 311-316.
64. Bergenholtz G, Lekholm U, Liljenberg B, Lindhe J (1983) Morphometric analysis of chronic inflammatory periapical lesions in root-filled teeth. *Oral Surg Oral Med Oral Pathol* 55: 295-301.
65. Kontiainen S, Ranta H, Lautenschlager I (1986) Cells infiltrating human periapical inflammatory lesions. *J Oral Pathol* 15: 544-546.
66. Yu SM, Stashenko P (1987) Identification of inflammatory cells in developing rat periapical lesions. *J Endod* 13: 535-540.
67. Anneroth G, Norberg KA (1968) Adrenergic vasoconstrictor innervation in the human dental pulp. *Acta Odontol Scand* 26: 89-93.
68. Pohto P (1972) Sympathetic adrenergic innervation of permanent teeth in the monkey (*Macaca*). *Acta Odontol Scand* 30: 117-126.
69. Brown AC, Barrow BL, Gadd GN, Van Hassel HJ (1969) Tooth pulp transcapillary osmotic pressure in the dog. *Arch Oral Biol* 14: 491-502.
70. Maita E, Simpson MD, Tao L, Pashley DH (1991) Fluid and protein flux across the pulp-dentine complex of the dog in vivo. *Arch Oral Biol* 36: 103-110.
71. Knutsson G1, Jontell M, Bergenholtz G (1994) Determination of plasma proteins in dentinal fluid from cavities prepared in healthy young human teeth. *Arch Oral Biol* 39: 185-190.
72. Vongsavan N, Matthews B (1992) Changes in pulpal blood flow and in fluid flow through dentine produced by autonomic and sensory nerve stimulation in the cat. *Proc Finn Dent Soc* 88: 491-497.
73. Vongsavan N, Matthews B (1992) Fluid flow through cat dentine in vivo. *Arch Oral Biol* 37:175-185.
74. Pashley DH (1985) Dentin-predentin complex and its permeability: physiologic overview. *J Dent Res* 64 Spec 613-620.
75. Bishop MA (1992) Extracellular fluid movement in the pulp; the pulp/dentin permeability barrier. *Proc Finn Dent Soc* 88: 331-335.
76. Lund T, Wiig H, Reed RK, Aukland K (1987) A "new" mechanism for oedema generation: strongly negative interstitial fluid pressure causes rapid fluid flow into thermally injured skin. *Acta Physiol Scand* 129: 433-435.

Copyright: © 2016 Silva. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Silva L (2016) A Literature Review of Inflammation and Its Relationship with the Oral Cavity. *Glob J Infect Dis Clin Res* 2(1): 001-007. DOI: 10.17352/2455-5363.000006